

# Severe Reaction to Bee Stings

## Emergency Plan of Action

Name: \_\_\_\_\_ D.O.B \_\_\_\_\_ Grade: \_\_\_\_\_

School: \_\_\_\_\_ Teacher: \_\_\_\_\_

If You See This	Do This
<ul style="list-style-type: none"> <li>• Itchy Skin</li> <li>• Swelling at sting site</li> <li>• Reported or suspected bee sting</li> </ul>	<ul style="list-style-type: none"> <li>• Stay with student</li> <li>• Stay calm. Keep student calm</li> <li>• Cleanse area with antiseptic</li> <li>• Observe for symptoms of anaphylaxis</li> </ul>
<ul style="list-style-type: none"> <li>• Hives spreading over body</li> <li>• Wheezing, difficulty talking, swallowing or breathing</li> <li>• Swelling of face, ears, lips or neck</li> <li>• Tingling/swelling of tongue</li> <li>• Vomiting</li> <li>• Extreme paleness/gray color, clammy skin</li> <li>• Loss of conscientiousness</li> </ul>	<ul style="list-style-type: none"> <li>• Administer EpiPen per orders.</li> <li>• Call 911 immediately</li> <li>• Do not leave student. Do not allow student to Stand up. Avoid any sudden movements after EpiPen has been administered.</li> <li>• Tell EMS that EpiPen was given. Provide used EpiPen.</li> </ul>
<p><b>Directions for use of EpiPen</b></p> <ul style="list-style-type: none"> <li>• Pull off gray or blue cap</li> <li>• Place black/orange tip against muscle on outer thigh, halfway between knee and hip</li> <li>• Press firmly until you hear a click</li> <li>• Hold in place for 10 seconds, then remove, massage area</li> <li>• Call 911</li> <li>• Keep student lying down. Do not allow student to get up</li> <li>• Do not return EpiPen to holder after use, give to EMS personnel or discard in sharps container</li> </ul>	
<p><b>After Administration</b>            Notify School nurse            Notify Parent            Complete Incident Report</p>	

\_\_\_\_\_  
Parent Signature

\_\_\_\_\_  
Date

\_\_\_\_\_  
Physician Name & Signature

\_\_\_\_\_  
Date

\*\*\* If your child has never had an anaphylactic reaction and does not need an EpiPen but you would like to keep a medication at his/her school to treat mild symptoms, please consult your pediatrician about the appropriate OTC medication. Remember all medications including OTC must be accompanied by a Medication Authorization Form in order to be administered in the school setting.